

String with noncommutative world-sheet and stringy instantons

I.V.Gorbunov* and A.A.Sharapov†

*Department of Physics, Tomsk State University,
Lenin Ave. 36, Tomsk, 634050 Russia*

hep-th/0201033

Abstract

The model of the bosonic string with the noncommutative world-sheet geometry is proposed in the framework of Fedosov's deformation quantization. The re-interpretation of the model in terms of bosonic string coupled to infinite multiplet of background fields is given and the link to the W -symmetry is discussed. For the case of $d=4$ Euclidean target space, the stringy counterparts of the Yang-Mills instantons are constructed in both commutative and noncommutative regimes.

PACS numbers: 11.25.-w, 04.60.Gw

Keywords: Deformation quantization, strings, instantons, noncommutative geometry.

1 Introduction

The passage from the paradigm of point particles to strings inevitably introduces a new fundamental scale related to the string size and leads to a new type of “fuzziness” in addition to the quantum-mechanical uncertainty. The quantum effects are proportional to the Planck constant $\hbar$, while the stringy effects are governed by the reciprocal of string tension α' and both of these parameters are responsible for a certain deformation of the classical phase-space and/or target-space geometry underlying the string theory. Although there is a strong evidence that varying these parameters one may interpolate between five known string models and even 11-dimensional supergravity, a constructive definition of the full theory in the “intermediate region”, called M-theory, remains a mystery. As to the $\hbar$ -deformation, we have the powerful machinery of modern quantization methods allowing to handle efficiently the problem in many cases. (It is the requirement of quantum consistency that appears to be a fundamental vehicle for

*

†

* *e-mail:* ivan@phys.tsu.ru

† *e-mail:* sharapov@phys.tsu.ru

identifying the five string theories as appropriate points on $(\hbar, \alpha')$ -scan of M-theory.) Meanwhile, the mathematical structure behind the α' -deformation is less well understood.

Nowadays most of the hope for obtaining a microscopic description of M-theory centers around the M(atr)ix theory [1, 2]. Perhaps, the simplest way to see how the Matrix models are related to string theory is to start with the “Poisson formulation” of the bosonic string [3, 4],

$$S = \int_M \omega (\{X^A, X^B\}^2 - \alpha^2) . \quad (1)$$

Here X^A are the target-space coordinates and $\{\cdot, \cdot\}$ stands for the Poisson bracket associated to the area two-form ω on the string world-sheet M . At the level of classical equations of motion the model (1) is equivalent to the Nambu-Goto string with $\alpha' = \alpha/2\pi$. The Matrix theory approach is based on exploiting a similarity between algebraic properties of the Poisson bracket and the matrix commutator. Considering the coordinates X^A to be Hermitian $N \times N$ -matrices and making a formal substitutions

$$\{\cdot, \cdot\} \Rightarrow i[\cdot, \cdot], \quad \int_M \omega \Rightarrow \text{Tr} , \quad (2)$$

in the action (1) one comes to what is known as IKKT matrix model¹ [3]:

$$S_{\text{IKKT}} = -\text{Tr} ([X^A, X^B]^2 + \alpha^2) . \quad (3)$$

Up to the constant term this action function can be viewed as the result of dimensional reduction to a point of $U(N)$ Yang-Mills theory. It is assumed that taking a certain large- N limit of this model one can approximate the bosonic string action (1) provided the eigenvalue distributions of the operators X^A are smooth enough. For two-torus and sphere such a limit can be traced explicitly, see e.g. [2].

It is very fruitful in physics to treat the operators as not to be given *a priori* but resulting from the quantization of phase-space observables. Assuming the IKKT model to be somehow related to the string theory it is more than natural to consider a quantization of the Poisson bracket entering to the action (1). As we will show below this can be actually done within the framework of Fedosov’s deformation quantization [5, 6]. What we get as a result may be regarded as the string with noncommutative world-sheet. From a general viewpoint this approach, as such, is very similar to that of noncommutative field theory, if not identical in two respects. First, the noncommutativity of geometry underlying our model is *dynamical*, i.e. governed by the unconstrained field ω , in contrast to a constant Poisson bracket used in the known models of noncommutative field theory. The second point, closely related to the first one, is that consistent definition of the star-product requires introducing a *symplectic connection* respecting form ω . Unlike the metric connection, the symplectic one is not completely determined in terms of the structure it preserves, i.e. via the form ω , and thereby it should be considered as an additional

¹More precisely, we consider only bosonic sector of this model.

dynamical field of the theory. The important observation behind our construction is that both prerequisites of the Fedosov quantization – symplectic structure and connection – are already contained in the string theory in a form of the Polyakov metric on the string world-sheet. Namely, we identify ω with the Riemannian volume-form, which is obviously preserved by the metric connection.

In this respect, the Deformation quantization allows us to link the Matrix and Noncommutative field theories in the quite natural way. The details of the construction are presented in the Section 2. There we also show that the model admits a straightforward re-interpretation as the conventional bosonic string coupled to a special background of external fields. This background is recognized as the one compatible with W -type symmetry generated by internal star-automorphisms of the theory. In so doing, the deformation parameter is naturally identified with the fundamental stringy scale α' .

Returning to the question about the nature of α' -deformation we may say that, in certain respects, it can be put in the same mathematical framework as $\hbar$ -deformation, i.e. in the framework of Deformation quantization theory.

In Section 3 we address the problem of constructing exact solutions to our model. On the whole, the situation here closely resembles the present state of the art of finding explicit solutions to the YM equations: for arbitrary dimension and signature the problem looks hopeless, but in special case of four-dimensional Euclidean space the series of instanton solutions can explicitly be constructed. Using the similarity between the forms of IKKT and YM action functionals we derive the stringy counterpart for the self-duality condition and then resolve it in a purely algebraic manner. In the commutative limit these solutions pass to the holomorphic curves in $\mathbb{C}^2 \sim \mathbb{R}^4$ which seem to exhaust all globally minimal two-surfaces in $\mathbb{R}^4$. It should be noted, however, that contrary to the case of YM instantons these are not localized on the world-sheet in both commutative and noncommutative regimes. The reason is that we are confined to solutions with analytical dependence on the deformation parameter. The existence of nonanalytical solutions remains an open question and we conclude with some remarks upon a possible line of attack on the problem.

2 Deformation of bosonic string

Consider the world-sheet of bosonic string as a Riemannian manifold M with a metric g_{ij} , $i, j = 1, 2$. Being a two-dimensional manifold, M also carries a symplectic structure given by the canonical volume form $\omega = \sqrt{g} du \wedge dv$, $g = |\det(g_{ij})|$. In fact, M is Kählerian manifold and both structures can be combined to one degenerate Hermitian form

$$\Lambda^{ij} = g^{ij} + \sqrt{-1}\omega^{ij}, \quad (4)$$

where g^{ij} and ω^{ij} are the inverse Riemannian and symplectic metrics, respectively. The right and left null-vectors of Λ can be visualized as those spanned light-cone at each point of M .

In the conventional string theory the coordinates of target space $X^A(\sigma, \tau)$ are considered to be bosonic fields forming a commutative algebra with respect to the ordinary pointwise multiplication of functions on M . The noncommutative generalization of the theory is obtained by deforming the algebra of classical fields. More precisely, we introduce a new associative operation, called star-product, being an one-parametric deformation of the pointwise one. This $*$ -product is made of Hermitian structure (4) and its first several terms are given by

$$\begin{aligned} X^A * X^B = & X^A \cdot X^B + \frac{\nu}{2} \Lambda^{ij} \nabla_i X^A \nabla_j X^B + \frac{\nu^2}{8} \Lambda^{i_1 j_1} \Lambda^{i_2 j_2} \nabla_{i_1} \nabla_{i_2} X^A \nabla_{j_1} \nabla_{j_2} X^B \left(1 - \frac{\nu}{4} R\right) + \\ & + \frac{\nu^3}{48} \Lambda^{i_1 j_1} \Lambda^{i_2 j_2} \Lambda^{i_3 j_3} \nabla_{i_1} \nabla_{i_2} \nabla_{i_3} X^A \nabla_{j_1} \nabla_{j_2} \nabla_{j_3} X^B + O(\nu^4), \end{aligned} \quad (5)$$

where ν is a deformation parameter, ∇_i and R are the covariant derivative and the scalar curvature associated to the metric g_{ij} . The higher terms $O(\nu^4)$ are restored step-by-step from an associativity condition for the $*$ -product. The explicit recurrent formulas can be found in [7, 8, 9].

Generally speaking, there are many other star-products associated to the same metric. For example, in Eq. (4) tensors ω and g can be combined with arbitrary complex coefficients. The above $*$ -product corresponds to a nonlinear generalization of the Wick symbol associated with the usual creation-annihilation operators. The use of the Wick star-product is seemed to be natural in the context of string theory since it implies, for example, the commutativity of left (right) modes of the closed string. More precisely, the product of functions $X^A(x^\pm)$ depending on one of curvilinear light-cone coordinates $x^\pm(u, v)$,

$$\Lambda^{ij} \partial_j x^+ = \partial_j x^- \Lambda^{ji} = 0, \quad (6)$$

does not deformed since $x^\pm * x^\pm = x^\pm \cdot x^\pm$ for the Wick $*$ -product (5).

To define the deformed string action we need one more ingredient of the deformation quantization – the trace functional on $*$ -algebra. It is defined as any linear functional vanishing on the star-commutator of functions

$$\begin{aligned} [X^A, X^B] &\equiv X^A * X^B - X^B * X^A = \\ &= \nu \{X^A, X^B\} + \frac{\nu^2}{2} \omega^{i_1 j_1} g^{i_2 j_2} \nabla_{i_1} \nabla_{i_2} X^A \nabla_{j_1} \nabla_{j_2} X^B \left(1 - \frac{\nu}{4} R\right) + \\ &+ \frac{\nu^3}{48} (\Lambda^{i_1 j_1} \Lambda^{i_2 j_2} \Lambda^{i_3 j_3} - \Lambda^{j_1 i_1} \Lambda^{j_2 i_2} \Lambda^{j_3 i_3}) \nabla_{i_1} \nabla_{i_2} \nabla_{i_3} X^A \nabla_{j_1} \nabla_{j_2} \nabla_{j_3} X^B + O(\nu^4). \end{aligned} \quad (7)$$

The trace functional is known to exist for any symplectic manifold and can be represented in the form

$$\text{Tr } f = \int_M d\mu f, \quad \text{Tr } [f, g] = 0, \quad (8)$$

where the trace-measure $d\mu$ is uniquely determined up to an overall constant. As is shown in Ref. [6], the measure has a form of series in ν with coefficients depending on the curvature tensor

and its derivatives. In our case this series starts with

$$d\mu = \omega \left(1 - \frac{\nu}{4} R + \frac{\nu^2}{24} \square R + O(\nu^3) \right), \quad (9)$$

$\square$ being the two-dimensional Laplace operator.

The action functional for the deformed string can now be written as

$$\begin{aligned} S_{def} &= \frac{1}{\nu^2} \int d\mu ([X^A, X^B] * [X_A, X_B] - \nu^2 \alpha^2) = \\ &= \int \omega \left(\{X^A, X^B\}^2 - \alpha^2 + \frac{\nu}{4} R \{X^A, X^B\}^2 + \nu \{X_A, \{X^B, X^B\}\} \square X_B + \right. \\ &\quad + \nu^2 \left[\frac{1}{12} \Lambda^{i_1 j_1} \Lambda^{i_2 j_2} \Lambda^{i_3 j_3} \nabla_{i_1} \nabla_{i_2} \nabla_{i_3} X^A \nabla_{j_1} \nabla_{j_2} \nabla_{j_3} X^B \{X_A, X_B\} + \right. \\ &\quad + \frac{1}{4} (\omega^{i_1 j_1} g^{i_2 j_2} \nabla_{i_1} \nabla_{i_2} X^A \nabla_{j_1} \nabla_{j_2} X^B)^2 - \frac{R}{8} \Lambda^{i_1 j_1} \Lambda^{i_2 j_2} \nabla_{i_1} \nabla_{i_2} X^A \nabla_{j_1} \nabla_{j_2} X^B \{X_A, X_B\} + \\ &\quad + \frac{1}{8} \Lambda^{i_1 j_1} \Lambda^{i_2 j_2} \nabla_{i_1} \nabla_{i_2} \{X^A, X^B\} \nabla_{j_1} \nabla_{j_2} \{X_A, X_B\} - \\ &\quad - \frac{1}{4} \Lambda^{i_1 j_1} \Lambda^{i_2 j_2} \nabla_{i_1} \nabla_{i_2} X^A \nabla_{j_1} \nabla_{j_2} X^B \square \{X_A, X_B\} + \\ &\quad \left. \left. - \frac{R}{8} \square \{X^A, X^B\} \{X_A, X_B\} + \frac{1}{24} \square R (\{X^A, X^B\}^2 - \alpha^2) \right] \right) + O(\nu^3). \end{aligned} \quad (10)$$

Besides the global Poincaré invariance, the action is invariant, by construction, under world-sheet diffeomorphisms

$$\delta X^A = \xi^i \partial_i X^A, \quad \delta g_{ij} = \xi^k \partial_k g_{ij} + g_{kj} \partial_i \xi^k + g_{ik} \partial_j \xi^k, \quad (11)$$

and internal *-automorphisms of fields X^A ,

$$\delta X^A = [\epsilon, X^A], \quad \delta g_{ij} = 0, \quad (12)$$

where ξ^i and ϵ are gauge parameters. In the commutative limit (i.e. when *-commutator reduces to the Poisson bracket) the gauge transformations (12) form a subgroup of the area-preserving diffeomorphisms (11). For this reason we refer to internal *-automorphisms (12) as a group of *quantized symplectomorphisms*. The latter is also known under the name of W_∞ -type symmetry and it has been intensively studied in the context of conformal field theory and integrable models [10, 11]. Thus, we see that although the commutative limit of our model is equivalent (at least classically) to the Polyakov string the gauge symmetries underlying both the formulations are quite different and, in a sense, complementary to each other: the conformal transformations of the metric rescale the symplectic structure, while the symplectomorphisms preserve it.

The variation of the action (10) leads to the following equations of motion:

$$-\frac{\nu^2}{4} \frac{\delta S}{\delta X^A} = [[X_A, X_B], X^B] = 0, \quad (13)$$

$$\begin{aligned}
\frac{1}{\sqrt{g}} \frac{\delta S}{\delta g_{kl}} = & -\frac{1}{2} g^{kl} (\{X^A, X^B\}^2 + \alpha^2) + \nu \left[-\frac{1}{8} g^{kl} R \{X^A, X^B\}^2 + \right. \\
& + \frac{1}{4} \omega^{ik} \omega^{jl} \nabla_i \nabla_j \{X^A, X^B\}^2 - 2 g^{ki} g^{lj} \{X_A, \{X^A, X^B\}\} \nabla_i \nabla_j X_B + \\
& + \frac{1}{2} g^{kl} \{X_A, \{X^A, X^B\}\} \square X_B + \frac{1}{2} g^{kl} \{X^A, X^B\} \{X_A, \square X_B\} - \\
& \left. - \frac{1}{2} (g^{ik} g^{jl} + g^{il} g^{jk} - g^{ij} g^{kl}) (\nabla_i \{X_A, \{X^A, X^B\}\}) \nabla_j X_B \right] + O(\nu^2) = 0.
\end{aligned} \tag{14}$$

By construction, these equations are known to be invariant under the $*$ -automorphisms, but only a part of them is written in a manifestly covariant form, namely, as double $*$ -commutator of X 's. Unfortunately, the compact algebraic representation is lacking for the variation with respect to metric. It is also seen that the lowest order of Eqs. (14) determines only the conformal factor $\sqrt{g}$, not the entire metric g_{ij} .

As in the case of undeformed string, the reparametrization invariance of the model (11) allows one to bring the metric to the conformally flat form,

$$g_{ij} = \sqrt{g} \eta_{ij}. \tag{15}$$

In this gauge the system of equations (14) breaks on two different parts. The trace part of the variation allows one to define the conformal factor $\sqrt{g}$ in a pure algebraic way, expressing it order-in-order via the fields X^A . Substitution of this expression to the traceless part of the system gives two constraints on X^A in addition to the equations of motion (13). It is the appearance of these constraints that drastically differs our model from its matrix counterpart (3). As it will be clear bellow (see Eqs. (17)-(19)) they have the same origin and interpretation as the Virasoro constraints for the conventional bosonic string coupled to background fields.

Remarkably, the system (13), (14) admits a special class of solutions being a deformation of the ordinary string dynamics,

$$\square X^A = O(\nu^2), \quad g_{ij} = \alpha^{-1} \partial_i X^A \partial_j X_A \left(1 + \frac{\nu}{4} R\right) + O(\nu^2), \tag{16}$$

where R and $\square$ are the curvature and Laplace operator associated to the *induced* world-sheet metric. The existence of such solutions is provided by the special form of the first “quantum correction” to the undeformed action functional: it is given by square of undeformed string equations of motion plus the curvature-depending term, which can be removed by rescaling of metric.

Let us comment upon the general structure of the deformed string action. In any order in ν it is given by fourth-order polynomials in the derivatives of the fields X^A contracted with the help of flat target-space metric. Identifying the deformation parameter ν with the reciprocal of string tension $\alpha' = \alpha/2\pi$ one may treat the deformation as the interaction vertices for the usual bosonic string moving in a very special Poincaré-invariant background – all the background fields are proportional to $\eta_{AB}\eta_{CD}$. It is not hard to guess about the most general action functional respecting the gauge transformations (11), (12) and leading to a richer content of background

fields. It reads

$$S = S_g + \int \mathcal{L}_*(X), \quad (17)$$

$$\mathcal{L}_*(X) = d\mu \sum_{m=0}^{\infty} L_{A_1 A_2 \dots A_m} X^{A_1} * X^{A_2} * \dots * X^{A_m},$$

where $L_{A_1 A_2 \dots A_m}$ are constant tensors and S_g is an arbitrary scalar functional of metric². Expanding the Lagrangian $\mathcal{L}_*$ in terms of the derivatives of fields X^A , we get

$$\mathcal{L}_*(X) = \omega \sum G_{A_1 \dots A_m}^{(f)}(X) \nabla_{i(n_1)} X^{A_1} \dots \nabla_{i(n_m)} X^{A_m} f^{i(n_1) \dots i(n_m)}(R, \nabla R, \dots), \quad (18)$$

$$\nabla_{i(n_k)} \equiv \nabla_{i_1} \nabla_{i_2} \dots \nabla_{i_{n_k}}.$$

Here tensors $G^{(f)}(X)$ play the role of the background fields compatible with the fundamental symmetries of the model and their structure is completely determined by the tensors L . Passing to the dimensionless coordinates, $X^A \rightarrow \sqrt{\alpha'} X^A$, we turn the expansion (18) to that in powers of α' or, what is the same, in a number of derivatives. The first several terms of the resulting expansion are

$$\mathcal{L}_* = \omega \left(\frac{1}{\alpha'} V(X) + g^{ij} \partial_i X^A \partial_j X^B G_{AB}(X) + \omega^{ij} \partial_i X^A \partial_j X^B H_{AB}(X) + D(X)R + O(\alpha') \right). \quad (19)$$

The background fields V, G, H, D associated naturally to the tachyon, metric, antisymmetric tensor and dilaton are not completely arbitrary – as they arise from the same set of tensors L – but subject to certain constraints. In particular,

$$D(X) = \frac{1}{4} V(X) = \sum_{k=0}^{\infty} L_{A_1 A_2 \dots A_k} X^{A_1} X^{A_2} \dots X^{A_k}. \quad (20)$$

It would be interesting to find the full system of the constraints ensuring the invariance of the above action with respect to $*$ -automorphisms and compare them with the known low-energy effective equations of motion for the background fields.

The above consideration allows one to treat the deformed string action as a special highly symmetrical phase of the conventional string theory rather than a completely new model.

3 Noncommutative string instantons and holomorphic curves

The form of the string action (10) is similar to that of (noncommutative) Yang-Mills theory and this analogy extends, in a sense, to their equations of motion: in both cases the immediate search for nontrivial exact solutions seems to be a hopeless task. The alternative appears, however, when one studies the Euclidean version of Yang-Mills theory in four dimensions. Here it is possible to present the Yang-Mills Lagrangian as the square of chiral part of the strength

²It is pertinent to note that, living aside S_g , the functional (17) is nothing but the action of the general multi-matrix model [12] written in terms of $*$ -product and the respective trace functional. This observation suggests another link to the W -symmetry and related integrable models [12, 13, 14, 15] although we will not dwell on this point here.

tensor and replace the initial equations of motion by stronger ones – (anti-)self-duality conditions. This is how the remarkable instanton solutions emerge. In this section we show that the same “instanton philosophy” works well in the case of commutative and noncommutative string theory providing us with class of easy-to-construct and study solutions.

We begin with the undeformed string action

$$S = \int \omega \left(\{X^A, X^B\}^2 + \frac{\alpha^2}{2} \right), \quad (21)$$

with $\mathbb{R}^4$ being a target space. Notice that the sign before the last term differs from that in Eq. (1) due to the Euclidian signature, otherwise the model has no classical solutions. The respective equations of motion read

$$\begin{aligned} \omega^{ij} \frac{\delta S}{\delta \omega^{ij}} &= \{X^A, X^B\} \{X_A, X_B\} - \frac{\alpha^2}{2} = 0 \\ -\frac{1}{4} \frac{\delta S}{\delta X^A} &= \{X^B, \{X_B, X_A\}\} = 0. \end{aligned} \quad (22)$$

Given the boundary conditions, the solutions to this system describe the minimal two-surfaces in $\mathbb{R}^4$ and a part of them can be obtained by the same trick one uses to derive the Yang-Mills instantons. Namely, we can rewrite the action (21) in the form

$$\begin{aligned} S &= \frac{1}{2} \int_M \omega (\{X^A, X^B\} + \alpha^{AB})_{\pm}^2 + \int_{\partial M} \theta, \\ \theta &= 2\alpha_{\pm}^{AB} X_A \partial_i X_B dl^i. \end{aligned} \quad (23)$$

Here subscripts $\pm$ stand for the (anti-)self-dual part of a second-rank skew-symmetric tensor on $\mathbb{R}^4$,

$$H_{\pm}^{AB} = H^{AB} \pm \frac{1}{2} \epsilon^{ABCD} H_{CD}, \quad (24)$$

$\alpha^{AB} = -\alpha^{BA}$ is a constant tensor normalized as $(\alpha_{\pm}^{AB})^2 = \alpha^2$, and dl^i is a line element on the boundary ∂M . Because of Euclidian signature we have the obvious inequality

$$S \geq \int_{\partial M} \theta, \quad (25)$$

and equality is attained iff the following conditions are satisfied:

$$(\{X^A, X^B\} + \alpha^{AB})_{\pm} = 0. \quad (26)$$

The solutions to these equations provide the minima for the action functional and may be thought of as string counterparts for the Yang-Mills instantons. The initial equations of motion (22) are obvious consequences of the obtained ones.

It is instructive to rewrite the system (26) in $\text{SO}(3)$ -covariant form

$$\{X^4, X^a\} = \frac{1}{2} \varepsilon_{abc} \{X^b, X^c\} - \alpha_{\pm}^{4a}, \quad a, b, c = 1, 2, 3. \quad (27)$$

Using the rotation symmetry we bring the vector α^{4a} to the form $(0, 0, \alpha)$. Now let us introduce the complex coordinates in $\mathbb{R}^4 = \mathbb{C}^2$ by the rule

$$Z = X_1 + iX_2, \quad W = X_3 + iX_4. \quad (28)$$

In these coordinates the self-duality equations (26) read

$$\{Z, W\} = 0, \quad \{Z, \bar{Z}\} + \{W, \bar{W}\} = 2i\alpha. \quad (29)$$

The first Poisson bracket means that fields $Z(u, v)$ and $W(u, v)$ are functionally dependent, say $W = f(Z)$. Then the second equation is just a definition of the induced volume form on the holomorphic curve $Z = f(W)$,

$$\omega = (2i\alpha)^{-1} (dZ \wedge d\bar{Z} + dW \wedge d\bar{W}) = \left(\frac{1 + |f'(Z)|^2}{2i\alpha} \right) dZ \wedge d\bar{Z}. \quad (30)$$

In such a way we reproduce the well-known fact from the geometry: any holomorphic surface in a Kähler manifold is globally minimal. So, the complex geometry delivers a simple method for generating exact solutions. For example, equation $Z^2 = W$ being resolved in a parametric form, gives

$$X_1 = u, \quad X_2 = v, \quad X_3 = u^2 - v^2, \quad X_4 = 2uv, \quad (31)$$

that defines (after compactification of $\mathbb{R}^4 = \mathbb{C}^2$ to $\mathbb{C}P^2$) the well-known minimal embedding of two-sphere $S^2 \sim \mathbb{C}P^1$ into $\mathbb{C}P^2$ as a projective line. The high-genus minimal surfaces arise as (hyper)elliptic curves $Z^2 = P(W)$, $P(W)$ being a polynomial without multiple roots.

The generalization of the above constructions to the noncommutative case is straightforward. Up to an inessential boundary term the action (10) can be rewritten as

$$S = \frac{1}{2} \int_M d\mu \left([X^A, X^B] + \alpha^{AB} \right)_\pm * \left([X_A, X_B] + \alpha_{AB} \right)_\pm. \quad (32)$$

Then the global minimum of the functional is achieved provided

$$([X^A, X^B] + \alpha^{AB})_\pm = 0. \quad (33)$$

Of course, the transition from (32) to (33) requires a justification as we deal now with the $*$ -square integrated with respect to the trace measure, and the both are not positive definite in general. The implication is really correct due to the following line of reasons. First, we confine ourselves by the fields to be formal power series in the deformation parameter ν with coefficients being smooth functions on a compact manifold M ,

$$X^A = X_0^A + \nu X_1^A + \nu^2 X_2^A + \dots, \quad (34)$$

$$g^{ij} = g_0^{ij} + \nu g_1^{ij} + \nu^2 g_2^{ij} + \dots. \quad (35)$$

Then, ignoring a subset of measure zero in M , we may assume the fields to be defined on a closed disc with the Darboux coordinates u, v , so that $\omega = du \wedge dv$. Since, locally, all the deformation

quantizations are known to be equivalent to each other there exist a formally invertible operator $B = 1 + \nu B_1 + \nu^2 B_2 + \dots$, B_k being finite-order differential operators, intertwining a given $*$ -product with the Weyl-Moyal one [6]. Explicitly,

$$f * g = B^{-1}(B(f) *_{WM} B(g)). \quad (36)$$

As this takes place, the trace of $*$ -product of two functions is given by the well-known Weyl's formula

$$Tr(f * g) = \int dudv B(f) \cdot B(g), \quad (37)$$

where $dudv = B^* d\mu$ is the canonical measure on the disc, and dot means ordinary multiplication. Now the implication (32) $\Rightarrow$ (33) immediately follows from nondegeneracy of the operator B .

Rewriting Eqs. (33) in terms of complex coordinates (28) we get

$$[Z, W] = 0, \quad [Z, \overline{Z}] - [W, \overline{W}] = 2\alpha\nu. \quad (38)$$

Proceeding in a close analogy with above analysis we can solve the first equation by assuming Z to be a $*$ -holomorphic function in W ,

$$Z = f_*(W) = \sum f_n(W^*)^n. \quad (39)$$

As to the second (real) equation, it serves then for the perturbative definition of the metric g_{ij} written in the conformal gauge (15).

It is pertinent to note that equations of the form (38) have also arose as those describing the static BPS membrane configurations in the BFSS matrix approximation [16, 17, 18]. The operator equations of the form (33) have been discussed recently in [19] in the context of noncommutative Yang-Mills instantons.

In principle, we may forget for the moment about the origin of Eqs. (38) and treat them as a set of commutation relations for abstract operators Z and W . From this viewpoint it is clear, that expressions (39) do not exhaust all possible solutions for the commutativity condition $[Z, W] = 0$: there are many commuting operators (and even matrices), which are not the functions of each other. It is very intuitive that solutions of such a type have no definite ‘‘classical limit’’ as $\nu \rightarrow 0$, and hence, poorly fit in a framework of the deformation quantization. This does not mean that such non-perturbative solutions have nothing to do with the noncommutative string paradigm at all. One of the possible scenario to reproduce them here is to extend the space of formal power series (34) to the class of the, so called, oscillating symbols having the form

$$f(u, v; \nu) \exp\left(\frac{i}{\nu} \varphi(u, v)\right), \quad (40)$$

where $f = f_0 + \nu f_1 + \dots$ is a formal power series in the deformation parameter and $\varphi, f_0, f_1, \dots$ are smooth functions on M . Expression (40) resembles the WKB-ansatz for the Wigner functions on the phase space and can be used to approximate the non-perturbative solutions for Eq. (38). It is these solutions, if any, that one may identify with *the genuine noncommutative string instantons*.

Acknowledgments

This work is supported in part by the INTAS under the grant N 00-262 and Russian Ministry of Education under the grant E-00-33-184. The work of I.V.G. is partially supported by RFBR under the grant N 00-02-17-956. The work of A.A.S. is partially supported by the RFBR Grant for Young Scientists N 01-02-06420.

References

- [1] T. Banks, W. Fischler, S. Shenker and L. Susskind, *M Theory as a Matrix Theory: A Conjecture*, Phys. Rev. **D55** (1997) P.5112-5128; hep-th/9610043.
- [2] W. Taylor, *The M(atrix) model of M-theory*, preprint MIT-CTP-2894; hep-th/0002016.
- [3] N. Ishibashi, H. Kawai, Y. Kitazawa and A. Tsuchiya, *A Large-N Reduced model as Superstring*, Nucl. Phys. **B498** (1997) P.467-491; hep-th/9612115.
- [4] A. Connes, M.R. Douglas and A. Schwarz, *Noncommutative Geometry and Matrix Theory: Compactification on Tori*, JHEP **9802** (1998) 003; hep-th/9711162.
- [5] B.V. Fedosov, *A Simple Geometric Construction of Deformation Quantization*, J. Diff. Geom. **40** (1994) P.213-238.
- [6] B.V. Fedosov, *Deformation quantization and Index Theory*, Akademie Verlag, Berlin, 1996.
- [7] M. Bordemann and S. Waldmann, *A Fedosov star product of Wick Type for Kähler Manifolds*. Lett. Math. Phys. **41** (1997) P.243-253; q-alg/9605012.
- [8] V.A. Dolgushev, S.L. Lyakhovich, A.A. Sharapov, *Wick type deformation quantization of Fedosov manifolds*, Nucl. Phys. **B606** [PM] (2001) P.647-672; hep-th/0101032.
- [9] A.V. Karabegov and M. Schlichenmaier, *Almost Kähler Deformation Quantization*, preprint math/0102169.
- [10] C.P. Pope, *Lectures on W-Algebras and W-Gravity*, Proceedings of 1991 Summer School in High Energy Physics and Cosmology, edited by E. Gava et al., World Scientific, 1992, P.827-867 and references therein; hep-th/9112076.
- [11] I. Bakas, *The Large-N Limit of Extended Conformal Symmetries*, Phys. Lett., **B228** (1989) P.57.
- [12] A. Morozov, *Integrability and Matrix Models*, Phys. Usp. **37** (1994) P.1-55; hep-th/9303139.
- [13] I.M. Gelfand and L.A. Dikii, *Asymptotic Behaviour of the Resolvent of Sturm-Liouville Equations and the Algebra of the Korteweg-de-Vries Equations*, Russ. Math. Surv. **30** (1975) P.77-113.
- [14] H. Awata, M. Fukuma, Y. Matsua and S. Odake, *Representation Theory of The $W_{1+\infty}$ Algebra*, Progr. Theor. Phys. Suppl. **118** (1995) P.343-374, and references therein; hep-th/9408158.
- [15] C. Castro, *W-Geometry from Fedosov Deformation Quantization*, J. Geom. Phys. **33** (2000) P.173-190; hep-th/9802023.
- [16] L. Cornalba and W. Taylor, *Holomorphic Curves from Matrices*, Nucl. Phys. **B536** (1998) P.513-552; hep-th/9807060.
- [17] L. Cornalba, *Matrix Representations of Holomorphic Curves on T_4* , JHEP 0008 (2000) 047; hep-th/9812184.
- [18] L. Cornalba and R. Schiappa, *Matrix Theory Star Products from the Born-Infeld Action*, Adv. Theor. Math. Phys. **4** (2000) P.249-269; hep-th/9907211.
- [19] A. Schwarz, *Noncommutative instantons: new approach*, Commun. Math. Phys. **221** (2001) P.433-450; hep-th/0102182.